

# Immunizations

## Age

### Vaccine

	Birth	1 month	2 months	4 months	6 months	12 months	15 months	18 months	19-23 months	2-3 years	4-6 years
<b>Hepatitis B</b>	<b>HepB</b>	<b>HepB</b>	see footnote 1	<b>HepB</b>							
<b>Rotavirus</b>		<b>Rota</b>	<b>Rota</b>	<b>Rota</b>							
<b>Diphtheria, Tetanus, Pertussis</b>		<b>DTaP</b>	<b>DTaP</b>	<b>DTaP</b>	see footnote 2	<b>DTaP</b>		<b>DTaP</b>			
<b>H. influenzae type b</b>		<b>Hib</b>	<b>Hib</b>	see footnote 3	<b>Hib</b>						
<b>Pneumococcal</b>		<b>PCV</b>	<b>PCV</b>	<b>PCV</b>	<b>PCV</b>		<b>PPV</b>				
<b>Inactivated Poliovirus</b>		<b>IPV</b>	<b>IPV</b>	<b>IPV</b>			<b>IPV</b>				
<b>Influenza</b>	<b>Influenza (Yearly)</b>										
<b>Measles, Mumps, Rubella</b>						<b>MMR</b>		<b>MMR</b>			
<b>Varicella</b>						<b>Varicella</b>		<b>Varicella</b>			
<b>Hepatitis A</b>						<b>HepA (2 Doses)</b>				<b>HepA Series</b>	
<b>Meningococcal</b>						<b>MCV4</b>					

  Range of recommended ages    
   Certain high risk groups    
   Catch-up immunization    
 Physicians are encouraged to use combination vaccines, when available, for childhood immunizations.

1. It is permissible to administer 4 doses of HepB when combination vaccines are administered after the birth dose. If monovalent HepB is used for doses after the birth dose, a dose at age 4 months is not needed.
2. The fourth dose of DTaP may be administered as early as age 12 months, provided 6 months have elapsed since the third dose. Administer the final dose in the series at age 4–6 years.
3. If PRP-OMP (PedvaxHIB® or ComVax® [Merck]) is administered at ages 2 and 4 months, a dose at age 6 months is not required. TriHibit® (DTaP/Hib) combination products should not be used for primary immunization but can be used as boosters after any Hib vaccine in children aged ≥12 months.



**BlueCross BlueShield of Texas**

*Experience. Wellness. Everywhere.™*

# Immunizations

## Age Group

Vaccine	7-10 years	11-12 years	13-18 years
Diphtheria, Tetanus, Pertussis	see footnote 1	Tdap	Tdap
Human Papillomavirus	see footnote 2	HPV (3 Doses)	HPV Series
Meningococcal	MCV4	MCV4	MCV4
Pneumococcal	PPV		
Influenza	Influenza (Yearly)		
Hepatitis A	HepA Series		
Hepatitis B	HepB Series		
Inactivated Poliovirus	IPV Series		
Measles, Mumps, Rubella	MMR Series		
Varicella	Varicella Series		

Range of recommended ages
  Certain high risk groups
  Catch-up immunization

Physicians are encouraged to use combination vaccines, when available, for childhood immunizations.

1. Minimum age: 10 years for BOOSTRIX® and 11 years for ADACEL™

2. Minimum age: 9 years.



**BlueCross BlueShield of Texas**

*Experience. Wellness. Everywhere.™*