

Client	HCSC STANDARD ALGORITHM <i>Implemented for Blue Cross Blue Shield of Texas</i>		
Measure Title	DIABETIC RETINAL EXAM		
Disease State	Diabetes, Retinopathy	Indicator Classification¹	Disease Management
Strength of Recommendation²	A		
Organizations Providing Recommendation	NCQA (HEDIS 2007 Technical Specification), The American Diabetes Association, The American Academy of Ophthalmology, The American College of Physicians		
Clinical Intent	To ensure that all members with diabetes receive a diabetic retinopathy screening test at a clinically appropriate frequency.		
Physician Specialties	Endocrinology, Family Practice, Geriatric Medicine, Internal Medicine		
Clinical Rationale	<p>Disease Burden</p> <ul style="list-style-type: none"> • Diabetes is a chronic, serious disease that affects approximately 14.7 million Americans. This disease is the leading cause of new cases of blindness among adults aged 20-74.[1] • After living with diabetes for 20 years, almost all patients with type 1 diabetes and 50 to 80 percent of those with type 2 diabetes will manifest signs of retinopathy.[2] • Retinopathy is a major cause of morbidity in patients with diabetes. The incidence of blindness, for example, is 25 times higher in patients with diabetes than in the general population. Furthermore, diabetic retinopathy is the most common cause of blindness in middle-aged subjects, accounting for at least 12 percent of all new cases in the United States each year.[3] <p>Reason for Indicated Intervention or Treatment</p> <ul style="list-style-type: none"> • Evidence supports that screening and early treatment for diabetic retinopathy is associated with a decreased rate of visual loss.[4-6] <p>Evidence supporting Intervention or Treatment</p> <ul style="list-style-type: none"> • In their cost-effectiveness analyses, Javitt and colleagues have reported that in patients with type 1 diabetes, annual screening (ophthalmoscopy with dilated pupils) for those without retinopathy and screening every six months for those with retinopathy followed by guideline concordant treatment would result in a saving of 70,000 to 80,000 person-years of sight and 60 to 80 million dollars annually in the United States.[7] In patients with type 2 diabetes, the same screening program and treatment would result in saving over 94,000 person-years of sight and over 250 million dollars per year.[8] • Appropriate screening and early detection of retinopathy preserves vision.[5, 9-11] • At least three randomized controlled trials have reported that photocoagulation for diabetic retinopathy preserves vision.[12-14] 		

Clinical Recommendations

The American Diabetes Association, the American Academy of Ophthalmology, and the American College of Physicians guidelines recommend[15-17] :

- Patients with type 1 diabetes (defined operationally as diabetes onset before age 30 years and requiring therapy with insulin) should have a complete examination by an ophthalmologist within three to five years after the onset of diabetes. Subsequent examinations should be determined by the presence and severity of retinopathy at the initial examination, but the minimum recommendation is annual examination. One exception is that screening is not indicated before puberty.[19]
- Patients with type 2 diabetes should have a complete examination by an ophthalmologist beginning at the time of diagnosis. Subsequent examinations or referral for treatment should be determined by the presence and severity of retinopathy at the initial examination, but the minimum recommendation is annual examination.[20]

Source	Adapted from Health Plan Employer Data and Information Set (HEDIS®) 2007 Technical Specification (unable to include evidence of negative retinal exam in year prior to measurement year as this is ascertained with result data that are not generally available via administrative claims). HBI's standard algorithm contains an additional exclusion for members with Type I diabetics. The numerator deviates from HEDIS in that the ophthalmology/optometry specific E & M codes for dilated eye exams do not require specialty codes.
Denominator	Continuously enrolled members ages 18 - 75 years by the end of the measurement year who were identified as having diabetes during the measurement year or year prior.
Denominator Exclusion	Members in the denominator with a diagnosis of polycystic ovaries at any time in the member's history who did not receive a diagnosis of diabetes during the measurement year or year prior, or members diagnosed with gestational diabetes, or steroid-induced diabetes during the measurement year or year prior.
Numerator	Members who received at least one screening exam for diabetic retinal disease by an eye care professional or had at least one office visit with an eye care professional during the measurement year.* <i>*Eye exams provided by eye care professionals are a proxy for dilated eye examinations because there is no administrative way to determine that a dilated exam was performed. Source: HEDIS 2007</i>
Interpretation of Score	High score implies better performance.
Physician Attribution	Score all physicians (in the selected specialties) who saw the member during the measurement year.

References

1. CDC. *National Diabetes Surveillance System*. 2004 [cited 2004 November 17th]; Available from: <http://www.cdc.gov/diabetes/statistics/prev/national/figpersons.htm>
2. Frank, R.N., *Diabetic retinopathy*. N Engl J Med, 2004. **350**(1): p. 48-58.
3. Klein, R., et al., *Association of ocular disease and mortality in a diabetic population*. Arch Ophthalmol, 1999. **117**(11): p. 1487-95.
4. Javitt, J.C. and L.P. Aiello, *Cost-effectiveness of detecting and treating diabetic retinopathy*. Ann Intern Med, 1996. **124**(1 Pt 2): p. 164-9.
5. Malone, J.I., et al., *Prevalence and significance of retinopathy in subjects with type 1 diabetes of less than 5 years' duration screened for the diabetes control and complications trial*. Diabetes Care, 2001. **24**(3): p. 522-6.
6. Chew, E.Y., et al., *The long-term effects of laser photocoagulation treatment in patients with diabetic retinopathy: the early treatment diabetic retinopathy follow-up study*. Ophthalmology, 2003. **110**(9): p. 1683-9.
7. Javitt, J.C., et al., *Detecting and treating retinopathy in patients with type I diabetes mellitus. A health policy model*. Ophthalmology, 1990. **97**(4): p. 483-94; discussion 494-5.
8. Javitt, J.C., et al., *Preventive eye care in people with diabetes is cost-saving to the federal government. Implications for health-care reform*. Diabetes Care, 1994. **17**(8): p. 909-17.
9. Singer, D.E., et al., *Screening for diabetic retinopathy*. Ann Intern Med, 1992. **116**(8): p. 660-71.
10. Fong, D.S., et al., *Retinopathy in diabetes*. Diabetes Care, 2004. **27 Suppl 1**: p. S84-7.
11. Zoega, G.M., et al., *Screening compliance and visual outcome in diabetes*. Acta Ophthalmol Scand, 2005. **83**(6): p. 687-690.
12. *Proliferative diabetic retinopathy: treatment with xenon-arc photocoagulation. Interim report of multicentre randomised controlled trial*. Br Med J, 1977. **1**(6063): p. 739-41.
13. *Photocoagulation for diabetic maculopathy. A randomized controlled clinical trial using the xenon arc. British Multicentre Study Group*. Diabetes, 1983. **32**(11): p. 1010-6.
14. *Photocoagulation for diabetic macular edema. Early Treatment Diabetic Retinopathy Study report number 1. Early Treatment Diabetic Retinopathy Study research group*. Arch Ophthalmol, 1985. **103**(12): p. 1796-806.
15. Aiello, L.e.a., *Retinopathy in Diabetes*. Diabetes Care, 2004. **27**(1).
16. *Screening guidelines for diabetic retinopathy. American College of Physicians, American Diabetes Association, and American Academy of Ophthalmology*. Ann Intern Med, 1992. **116**(8): p. 683-5.
17. ADA, *Standards of Medical Care in Diabetes* Diabetes Care, 2005. **28** ((Supplement 1): S4).
18. AAO, *Diabetic Retinopathy, in preferred practice pattern*. 2003, American Academy of Ophthalmology.
19. Klein, R., et al., *The Wisconsin epidemiologic study of diabetic retinopathy. II. Prevalence and risk of diabetic retinopathy when age at diagnosis is less than 30 years*. Arch Ophthalmol, 1984. **102**(4): p. 520-6.
20. Klein, R., et al., *The Wisconsin epidemiologic study of diabetic retinopathy. III. Prevalence and risk of diabetic retinopathy when age at diagnosis is 30 or more years*. Arch Ophthalmol, 1984. **102**(4): p. 527-32.

¹ **Indicator Classification** (Adapted from Health Plan Employer Data Information Set (HEDIS®) technical specifications)

Diagnosis	Measures applicable to patients receiving diagnostic workups for a symptom or condition that delineate appropriate laboratory or radiological testing to be performed (e.g. evaluation of thyroid nodule; pregnancy test in patients with vaginal bleeding or abdominal pain)
Effectiveness of Care	
Prevention	Measures applicable to asymptomatic individuals that are designed to prevent the onset of the targeted condition (e.g. immunizations).
Screening	Measures applicable to asymptomatic patients who have risk factors or pre-clinical disease, but in whom the condition has not become clinically apparent (e.g. pap smears; screening for elevated blood pressure).
Disease Management	Measures applicable to individuals diagnosed with a condition that are part of the treatment or management of the condition (e.g. cholesterol reduction in patients with diabetes; radiation therapy following breast conserving surgery; appropriate follow-up after acute event).
Medication Monitoring	Measures applicable to patients taking medications with narrow therapeutic windows and / or potential preventable significant side effects or adverse reactions (e.g. thyroid stimulating hormone (TSH) testing after levothyroxine dose change; hepatic enzyme monitoring for patients using antimycotic pharmacotherapy)
Medication Adherence	Measures applicable to patients taking medications for chronic conditions that are designed to assess patient adherence to medication (e.g. adherence to lipid lowering medication).
Utilization	Measures applicable to patients receiving treatment for a symptom or condition that advocate appropriate utilization of laboratory and pharmaceutical resources (e.g. conservative use of imaging for low back pain; inappropriate use of antibiotics for viral upper respiratory infection).

² Strength of Recommendation

Strength of Recommendation Based on a Body of Evidence




FIGURE 2. Algorithm for determining the strength of a recommendation based on a body of evidence (applies to clinical recommendations regarding diagnosis, treatment, prevention, or screening). While this algorithm provides a general guideline, authors and editors may adjust the strength of recommendation based on the benefits, harms, and costs of the intervention being recommended. (USPSTF = U.S. Preventive Services Task Force)