



CLINICAL PAYMENT AND CODING POLICY

If a conflict arises between a Clinical Payment and Coding Policy (CPCP) and any plan document under which a member is entitled to Covered Services, the plan document will govern. If a conflict arises between a CPCP and any provider contract pursuant to which a provider participates in and/or provides Covered Services to eligible member(s) and/or plans, the provider contract will govern. "Plan documents" include, but are not limited to, Certificates of Health Care Benefits, benefit booklets, Summary Plan Descriptions, and other coverage documents. BCBSTX may use reasonable discretion interpreting and applying this policy to services being delivered in a particular case. BCBSTX has full and final discretionary authority for their interpretation and application to the extent provided under any applicable plan documents.

Providers are responsible for submission of accurate documentation of services performed. Providers are expected to submit claims for services rendered using valid code combinations from Health Insurance Portability and Accountability Act (HIPAA) approved code sets. Claims should be coded appropriately according to industry standard coding guidelines including, but not limited to: Uniform Billing (UB) Editor, American Medical Association (AMA), Current Procedural Terminology (CPT®), CPT® Assistant, Healthcare Common Procedure Coding System (HCPCS), ICD-10 CM and PCS, National Drug Codes (NDC), Diagnosis Related Group (DRG) guidelines, Centers for Medicare and Medicaid Services (CMS) National Correct Coding Initiative (NCCI) Policy Manual, CCI table edits and other CMS guidelines.

Claims are subject to the code edit protocols for services/procedures billed. Claim submissions are subject to claim review including but not limited to, any terms of benefit coverage, provider contract language, medical policies, clinical payment and coding policies as well as coding software logic. Upon request, the provider is urged to submit any additional documentation.

Vectra DA Blood Test for Rheumatoid Arthritis

Policy Number: CPCPLAB039

Version 1.0

Enterprise Medical Policy Committee Approval Date: 1/25/2022

Plan Effective Date: May 1, 2022

Description

BCBSTX has implemented certain lab management reimbursement criteria. Not all requirements apply to each product. Providers are urged to review Plan documents for eligible coverage for services rendered.

Reimbursement Information:

The use of a multi-biomarker disease activity score for rheumatoid arthritis (e.g., Vectra DA score) **is not reimbursable**.



This policy does not pertain to general inflammation; for guidance on general inflammation testing, including erythrocyte sedimentation rate (ESR) and C-reactive protein (CRP), please see policy CPCPLAB49 General Inflammation Testing and policy CPCPLAB11 ANA/ENA Testing.

Procedure Codes

Codes
81490

References:

- Anderson, J., Caplan, L., Yazdany, J., Robbins, M. L., Neogi, T., Michaud, K., . . . Kazi, S. (2012). Rheumatoid arthritis disease activity measures: American College of Rheumatology recommendations for use in clinical practice. *Arthritis Care Res (Hoboken)*, 64(5), 640-647. doi:10.1002/acr.21649
- Bakker, M. F., Cavet, G., Jacobs, J. W., Bijlsma, J. W., Haney, D. J., Shen, Y., . . . Welsing, P. M. (2012). Performance of a multi-biomarker score measuring rheumatoid arthritis disease activity in the CAMERA tight control study. *Ann Rheum Dis*, 71(10), 1692-1697. doi:10.1136/annrheumdis-2011-200963
- Bakker, M. F., Jacobs, J. W., Verstappen, S. M., & Bijlsma, J. W. (2007). Tight control in the treatment of rheumatoid arthritis: efficacy and feasibility. *Ann Rheum Dis*, 66 Suppl 3, iii56-60. doi:10.1136/ard.2007.078360
- Brahe, C. H., Ostergaard, M., Johansen, J. S., Defranoux, N., Wang, X., Bolce, R., . . . Hetland, M. L. (2019). Predictive value of a multi-biomarker disease activity score for clinical remission and radiographic progression in patients with early rheumatoid arthritis: a post-hoc study of the OPERA trial. *Scand J Rheumatol*, 48(1), 9-16. doi:10.1080/03009742.2018.1464206
- Centola, M., Cavet, G., Shen, Y., Ramanujan, S., Knowlton, N., Swan, K. A., . . . Curtis, J. R. (2013). Development of a Multi-Biomarker Disease Activity Test for Rheumatoid Arthritis. *PLoS One*, 8(4). doi:10.1371/journal.pone.0060635
- Combe, B., Landewe, R., Daien, C. I., Hua, C., Aletaha, D., Alvaro-Gracia, J. M., . . . van Vollenhoven, R. (2017). 2016 update of the EULAR recommendations for the management of early arthritis. *Ann Rheum Dis*, 76(6), 948-959. doi:10.1136/annrheumdis-2016-210602
- Curtis, J. R., Flake, D. D., Weinblatt, M. E., Shadick, N. A., Ostergaard, M., Hetland, M. L., . . . Lanchbury, J. S. (2019). Adjustment of the multi-biomarker disease activity score to account for age, sex and adiposity in patients with rheumatoid arthritis. *Rheumatology (Oxford)*, 58(5), 874-883. doi:10.1093/rheumatology/key367
- Curtis, J. R., Greenberg, J. D., Harrold, L. R., Kremer, J. M., & Palmer, J. L. (2018). Influence of obesity, age, and comorbidities on the multi-biomarker disease activity test in rheumatoid arthritis. *Semin Arthritis Rheum*, 47(4), 472-477. doi:10.1016/j.semarthrit.2017.07.010

- Curtis, J. R., van der Helm-van Mil, A. H., Knevel, R., Huizinga, T. W., Haney, D. J., Shen, Y., . . . Weinblatt, M. E. (2012). Validation of a novel multibiomarker test to assess rheumatoid arthritis disease activity. *Arthritis Care Res (Hoboken)*, 64(12), 1794-1803. doi:10.1002/acr.21767
- England, B. R., Tiong, B. K., Bergman, M. J., Curtis, J. R., Kazi, S., Mikuls, T. R., . . . Michaud, K. (2019). 2019 Update of the American College of Rheumatology Recommended Rheumatoid Arthritis Disease Activity Measures. *Arthritis Care Res (Hoboken)*, 71(12), 1540-1555. doi:10.1002/acr.24042
- Firestein, G. (2020). Pathogenesis of rheumatoid arthritis - UpToDate. In P. L. Romain (Ed.), UpToDate. Retrieved from https://www.uptodate.com/contents/pathogenesis-of-rheumatoid-arthritis?source=search_result&search=ra&selectedTitle=6~150
- Hirata, S., Dirven, L., Shen, Y., Centola, M., Cavet, G., Lems, W. F., . . . Allaart, C. F. (2013). A multi-biomarker score measures rheumatoid arthritis disease activity in the BeSt study. *Rheumatology (Oxford)*, 52(7), 1202-1207. doi:10.1093/rheumatology/kes362
- Li, W., Sasso, E. H., Emerling, D., Cavet, G., & Ford, K. (2013). Impact of a multi-biomarker disease activity test on rheumatoid arthritis treatment decisions and therapy use. *Curr Med Res Opin*, 29(1), 85-92. doi:10.1185/03007995.2012.753042
- Li, W., Sasso, E. H., van der Helm-van Mil, A. H., & Huizinga, T. W. (2016). Relationship of multi-biomarker disease activity score and other risk factors with radiographic progression in an observational study of patients with rheumatoid arthritis. *Rheumatology (Oxford)*, 55(2), 357-366. doi:10.1093/rheumatology/kev341
- Ma, M. H. Y., Defranoux, N., Li, W., Sasso, E. H., Ibrahim, F., Scott, D. L., & Cope, A. P. (2020). A multi-biomarker disease activity score can predict sustained remission in rheumatoid arthritis. *Arthritis Res Ther*, 22(1), 158. doi:10.1186/s13075-020-02240-w
- Mc Ardle, A., Flatley, B., Pennington, S. R., & FitzGerald, O. (2015). Early biomarkers of joint damage in rheumatoid and psoriatic arthritis. *Arthritis Res Ther*, 17(1), 141. doi:10.1186/s13075-015-0652-z
- Mease, P. J. (2010). Improving the routine management of rheumatoid arthritis: the value of tight control. *J Rheumatol*, 37(8), 1570-1578. doi:10.3899/jrheum.091064
- NICE. (2019). Therapeutic monitoring of TNF-alpha inhibitors in rheumatoid arthritis. Retrieved from <https://www.nice.org.uk/guidance/dg36/chapter/1-Recommendations>
- NICE. (2020). Rheumatoid arthritis in over 16s. Retrieved from <https://www.nice.org.uk/guidance/qs33>
- Shmerling, R. (2020). Origin and utility of measurement of rheumatoid factors. Retrieved from https://www.uptodate.com/contents/origin-and-utility-of-measurement-of-rheumatoid-factors?topicRef=7503&source=see_link
- Singh, J. A., Saag, K. G., Bridges, S. L., Jr., Akl, E. A., Bannuru, R. R., Sullivan, M. C., . . . McAlindon, T. (2015). 2015 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. *Arthritis Care Res (Hoboken)*, 68(1), 1-25. doi:10.1002/acr.22783



Smolen, J. S., Breedveld, F. C., Burmester, G. R., Bykerk, V., Dougados, M., Emery, P., . . . van der Heijde, D. (2016). Treating rheumatoid arthritis to target: 2014 update of the recommendations of an international task force. *Ann Rheum Dis*, 75(1), 3-15. doi:10.1136/annrheumdis-2015-207524

Smolen, J. S., Landewé, R. B. M., Bijlsma, J. W. J., Burmester, G. R., Dougados, M., Kerschbaumer, A., . . . van der Heijde, D. (2020). EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2019 update. *Ann Rheum Dis*, 79(6), 685-699. doi:10.1136/annrheumdis-2019-216655

Taylor, P., & Maini, R. (2020). Investigational biologic markers in the diagnosis and assessment of rheumatoid arthritis. In J. O'Dell (Ed.), UpToDate. Retrieved from <https://www.uptodate.com/contents/investigational-biologic-markers-in-the-diagnosis-and-assessment-of-rheumatoid-arthritis>

van der Helm-van Mil, A. H. M., Knevel, R., Cavet, G., Huizinga, T. W. J., & Haney, D. J. (2013). An evaluation of molecular and clinical remission in rheumatoid arthritis by assessing radiographic progression. In *Rheumatology (Oxford)* (Vol. 52, pp. 839-846).

Venables, P., Maini, Ravinder. (2019). Clinical manifestations of rheumatoid arthritis. Retrieved from https://www.uptodate.com/contents/clinical-manifestations-of-rheumatoid-arthritis?search=Clinical%20manifestations%20of%20rheumatoid%20arthritis&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1

Policy Update History:

5/1/2022	New policy
----------	------------